

Effects of Air Pollution Related Respiratory Symptoms in Schoolchildren in Industrial Areas Rayong, Thailand

Paradee Asa^a and Wanida Jinsart^b

^a *Interdepartment of Environmental Science, Graduate School, Chulalongkorn University, Thailand*

^b *Department of Environmental Science, Faculty of Science, Chulalongkorn University, Thailand*

Abstract

The chronic effects of air pollution in school children living near industrial sites were investigated. There respiratory symptoms of 806 school children aged 9-12 years were examined by the American Thoracic Society's Division of Lung Diseases (ATS-DLD-78-C) questionnaire during February-August 2013. The selected elementary schools in this survey was based on the distance from Map Ta Phut Industrial Estate, area A within 1 km., area B 5 km. and area C 10 km. Logistic regression techniques were used to assess the association between prevalence of respiratory symptoms and independent variables. The average 24 hrs PM₁₀ and VOCs concentrations from 2011-2014 in area A were significantly higher than in area C ($p < 0.05$). Relatively, the prevalence of respiratory symptoms in area A were high with odds ratios (OR) = 3.41, (95% confidence intervals (CI) = 1.70-6.85) and in area B with OR = 1.36, (95% CI = 0.54-3.45), in comparison to area C. The prevalent of non-specific respiratory diseases (NSRD) and Persistent Cough and Phlegm (PCP) in boy student were higher than girls students with OR=2.17, (95% CI=1.33-3.53), whereas those factors such as age, residential years, home size, parental smoking habits, use of air conditioners and domestic pets were not associated. Exposure to particulate matter and volatile organic compounds arising from Industrial sites was associated with worse respiratory impairments in children.

Keywords: air pollution; respiratory symptoms; industrial area; children health

1. Introduction

Industry related air pollutants exert its adverse impacts on human health, particularly respiratory system. Exposure to PM₁₀ could effect on breathing, damage lung tissue, cause lung cancer and premature death (EPA, 1995). High level of Volatile organic compounds (VOCs) associated to increasing of asthma and high levels of the inflammatory marker nitric oxide in the exhaled breath in children (Rumchev *et al.*, 2002; 2004). Toxic VOCs, 1,3 butadiene, 1,2 dichloroethane and benzene were classified as carcinogenic in human (EPA, 2000).

Previous epidemiological studies have suggested relatively consistent associations between ambient air pollutions and human health (Pascal *et al.*, 2013). To illustrate, its effects on respiratory symptoms from traffic related air pollutions were reported in school children (Langkulsen *et al.*, 2006) or traffic policemen (Tamura *et al.*, 2003; Karita *et al.*, 2004); health effects caused by PM_{2.5}, PM₁₀, SO₂ (Aekplakorn *et al.*, 2003; Peled *et al.*, 2005) and NO_x from power plant were also reported by Yogev-Baggio *et al.* (2010). Furthermore, respiratory symptoms among residents exposed to

petrochemical pollution were previously reported in China (Wilson *et al.*, 2008) and in Argentina (Wichmann *et al.*, 2009).

Our study area, Map Ta Phut Municipality covered 165.575 km², 38 communities around the Map Ta Phut Industrial Estate (MTPIE) and 58,245 registered populations (MTP Municipality, 2014). The Map Ta Phut area is influenced by the sea wind which sweeps from the southwest to the northeast with a wind speed below 6ms⁻¹ (Pimpisut *et al.*, 2005). Many heavy industries and Petrochemical related manufactures were located in MTPIE, Rayong province and high level of air pollutions such as PM₁₀ and VOCs in this area were recorded (www.aqnis.pcd.go.th, 2015). Comparison of data from each monitoring site indicated that PM₁₀ and VOCs across sites were different from their major species and their concentrations which might be influenced by nearest potential emission sources. These conditions prompted us to investigate the extent of respiratory symptoms and their potential risk factors. Hence, the objective of this study was to assess the respiratory health effects on children exposed to PM₁₀ and VOCs living in the different distance from industrial sites by the epidemiological cross-sectional method.

2. Materials and Methods

2.1. Study area and population

The study areas were selected based on the ambient air quality levels and the distance from industrial sources in MTPIE. Fig. 1 illustrated the three areas, distanced from MTPIE, 1 km.(A), 5 km.(B) and 10 km.(C). The ambient air pollution levels in these studied areas were compared in Table 1 and 2. Four selected schools located in area A, two schools in area B and another two schools in area C respectively (Fig. 1). Schoolchildren aged 9-12 from these eight governmental primary schools volunteered to involve in this survey were 2,145.

2.2. Air pollution monitoring

Air quality data from 2011 to 2014 were obtained from the Air Quality and Noise Management Bureau (AQNIS), Pollution Control Department (PCD), Thailand (www.aqnis.pcd.go.th, 2015) and Map Ta Phut Industrial estate (MTPIE, 2014). Air pollution concentrations were measured following the standard methods, PM₁₀ by Beta attenuator air sampling and volatile organic compounds (VOCs) were collected using canisters and were analyzed by gas chromatography/mass spectrophotometer (GC/MS) following the US.EPA TO15 procedure. Composite samples taken

over a 24-h period were collected monthly. The air quality monitoring stations were located within the studied areas as shown in Fig. 1. The 24 hour average concentrations of PM₁₀, in 2011-2013 from station Mc, Nf, Tk, 29T, 74T and 30T were compared in Table 1. VOCs concentrations in 2012-2014 were obtained from MC, NF, TK, MT and NJ monitoring stations, Table 2.

2.3. Respiratory Questionnaires

American Thoracic Society's Division of Lung Diseases (ATS-DLD-78-C) questionnaire is a standard questionnaire using in NSRD and respiratory symptom study. The questionnaire consists of general information (13 items), respiratory symptoms (20 items), and family history (7 items) as details in (ATS-DLD, 2015). The analysis of the questionnaires was followed the epidemiology standard method, Ferris, 1978. The criteria of analysis were summarized in Table 3. Questionnaires randomly administered to 1500 students from total 2145 students in eight schools. The number of delivered questionnaires was higher than 10% precision level of sampling size, recommended 525 (Yamane, 1973). The survey was conducted during February -August 2013. This study was approved by Ethics Review Committee for Research Involving Human Research Subject, Health Science Group, Chulalongkorn University.




Figure 1. Study sites distance further from petrochemical industries: ----- area A within 1 km, area B 5 km and area C 10 km ★ VOCs monitoring stations: I = MC, II = NF, III = TK, IV = MT, V = NJ ● PM₁₀ monitoring stations: Mc, Nf, Tk, 29T, 74T, 30T ● Elementary schools: area A (1 = Wat Map-cha-lood, 2 = Ban Nong-fab, 3 = Wat Ta-kurn, 4 = Ban Map-ta-phut), area B (5 = Wat Khot-hin-mit-tra-phap, 6 = Wat Trok-yay-cha) and area C (7 = Wat Nong-snom, 8 = Wat Nuen-phra)

Table 1. Area characteristics, distance from MTPIE and 24 hrs average concentration of PM₁₀, in 2011-2013

Area	Characteristic	PM ₁₀ (µg/m ³)			P
		Range	Average	Frequency of exceeding standard	
A	Surrounded area within 1 km.				A VS B ns
	Station Mc	0-3471	<u>38.81</u>	-	A VS C *
	Station Nf	0-668	35.00	-	
	Station Tk	0-578	40.68	-	
	Station 29T	5-209	44.85	24/885 (2.7%)	
	Average	-	39.84	-	
B	Residential area within 5 km.				
	Station 74T	<u>11-107</u>	35.55	0/725 (0%)	B VS C ns
C	Residential area within 10 km.				-
	Station 30T	4.4-107	27.45	3/928 (0.3%)	

* p < 0.05 by one-way ANOVA

Data from the www.aqnis.pcd.go.th (2015) and MTPIE (2014)

2.4. Data analysis

A nonparametric method (Mann-Whitney U-test) was used to compare the levels of VOCs. Differences in the health-related parameters among the areas A, B and control area C were compared using Yates' chi-squared test. Logistic regression techniques were used to assess the association between prevalence of respiratory symptoms and independent variables such as gender, age, residential years, residential areas,

home size, family members, parental smoking habits, use of air conditioners and having domestic pets. The odds ratios (ORs) and 95% confidence intervals (CIs) were obtained as the outcome variables and precision weighting was applied to estimate the degree of association in this study. All statistical analyses were performed using the Statistical Package for Social Science (IBM SPSS Statistics 19; SPSS Inc, Chicago, IL, USA, 2010).

Table 2. Annual average concentration of VOCs in 2012-2014: Industrial Area (A) and Reference area (C)

Area	Air monitoring station	VOCs (µg/m ³)									
		VinylChloride	1,3 Butadiene	Dichloromethane	Chloroform	1,2 Dichloroethane	Benzene	Trichloroethylene	1,2 Dichloropropane	Tetrachloroethylene	Total VOCs
A	MC	0.21	0.18	0.97	0.16	0.35	1.23	0.27	0.11	0.14	3.61
	NF	0.06	0.29	0.66	0.29	0.30	1.37	0.16	0.23	0.13	3.49
	TK	0.18	<u>0.86*</u>	0.92	0.12	0.28	<u>2.33</u>	0.17	0.11	0.14	5.11*
	MT	0.93	<u>0.69*</u>	1.00	0.15	<u>1.21</u>	<u>2.60</u>	0.18	0.11	0.16	7.03*
	Total	0.35	<u>0.51</u>	0.89	0.18	<u>0.54</u>	<u>1.88</u>	0.78	0.14	0.14	4.81
C	NJ	0.05	0.05	0.41	0.12	0.20	<u>1.90</u>	0.17	0.10	0.14	3.13
Annual avg. standard		10	0.33	22	0.43	0.4	1.7	23	4	200	

*p < 0.05, Mann-Whitney U test,

The underline values are exceed the National Ambient Air Quality Standard (NAAQS),

Data from www.aqnis.pcd.go.th (2015)

Table 3. Definition of respiratory symptoms in ATS-DLD-78-C Questionnaire

Respiratory symptom		Definition
Non-Specific Respiratory Diseases: NSRD	Chronic Bronchitis (CB)	phlegm production from the chest two times/day for 4 days/week for 3 months/year for at least 3 years
	Bronchial Asthma (BA)	doctor diagnosed asthma and still have asthma
	Dyspnea and Wheezing (D&W)	wheezing or whistling in the chest apart from cold on most days or nights
Persistent Cough and Phlegm: PCP	Persistent Cough	cough apart from cold on most days more than 4 days/week for 3 months/year for at least 1 year
	Persistent Phlegm	congested in the chest or bring up phlegm, sputum, or mucus apart from cold on most days more than 4 days/week for 3 months/year for at least 1 year

3. Results and Discussion

3.1. Air quality and health effects

The air quality data in Map Ta Phut area were obtained from Pollution Control Department (PCD), Thailand. The VOCs and criteria air pollutants PM₁₀, SO₂, NO₂, CO and O₃ were recorded. The concentrations were different across monitoring sites. In this study we investigated the effect from two major pollutants, PM₁₀ and VOCs since the other gaseous pollutants average concentrations were not exceeded the NAAQS. PM₁₀ and VOCs in area A were found higher than that of area B and C (Table 1 and 2). Particularly, concentrations of PM₁₀ in area A were 2.7% frequency exceeded the standard level (Table 1) and three carcinogenic VOCs namely 1,3 butadiene, 1,2 dichloroethane and benzene were exceeded the NAAQS (Table 2). These could

associate with the high prevalence of respiratory symptoms in area A, B and C, 18 %, 8% and 5% respectively, $p < 0.05$, Table 4. Details of NSRD and PCP prevalence in each school and each area were included in Table 5. Total PCP prevalence in area A were higher than in C significantly. Our results indicated the association between respiratory prevalence in children and air pollutions which was in the same trend as previous studies in adult and pregnant subjects (Tanyanont *et al.*, 2012; Kongtip *et al.*, 2013; Phatrabuddha *et al.*, 2013). In addition, Thepanondh *et al.* (2011) reported that high concentration of benzene and 1,3 butadiene were detected at the stations near factories and the main roads, 1,2 dichloroethane observed near monitored site close to the plastic factories. Potentially, both stationary and mobile sources contributed to high level of VOCs.

Table 4. Air pollution in 2011-2014 and respiratory symptoms in schoolchildren (n=806)

Pollutant/ respiratory Symptoms	Area			(A and C) <i>p-value</i>
	A	B	C	
PM-10 (mean, µg/m ³)	39.84	35.55	27.45	< 0.05
VOCs (mean, µg/m ³)				
1,3 butadiene	0.51	-	0.05	
1,2 dichloroethane	0.54	-	0.20	
benzene	1.88	-	1.90	
total VOCs	4.81	-	3.13	
Prevalence (%)				
- Persistent Cough	4.5	2.3	0.8	< 0.05
- Persistent Phlegm	3.3	0.7	0	< 0.05
Any one of the respiratory Symptoms				
-Individual basis	58(14)	9(7)	12(5)	< 0.05
-Symptoms basis	77(18)	11(8)	12(5)	< 0.05
History of allergic diseases, n (%)				
Dust and Other chemical	96(22.8)	15(11.4)	47(18.5)	
Past illness, n (%)				
Respiratory Symptoms	116(27.6)	31(23.5)	40(15.7)	

Table 5. Number and prevalence (%) of respiratory symptoms in school children (n=806)

Area ^a	School	NSRD			PCP		Any one of the respiratory symptoms		
		CB	BA	D and W	Persistent Cough	Persistent phlegm	Both PCP	Individual basis	Symptoms basis
A n=420	MC	2(1.9)	7(6.6)	10(9.4)	9(8.5)	6(5.7)	2(1.9)	24(23)	34(32)
	NF	0	2(3.6)	6(10.7)	2(3.6)	3(5.3)	1(1.8)	10(18)	13(23)
	TK	0	3(3.6)	3(3.6)	0	2(2.4)	0	7(8)	8(10)
	MT	2(1.1)	4(2.3)	5(2.8)	8(4.6)	3(1.7)	1(0.6)	17(10)	22(13)
	total	4(0.9)	16(3.8)	24(5.7)	19(4.5)*	14(3.3)*	4(0.9)	58(14)*	77(18)*
B n=132	KH	1(1.7)	2(3.3)	2(3.3)	2(3.3)	1(0.7)	1(1.7)	6(10)	8(13)
	YC	0	2(2.8)	0	1(1.4)	0	0	3(4)	3(4)
	total	1(0.7)	4(3.0)	2(1.5)	3(2.3)	1(0.7)	1(0.7)	9(7)	11(8)
C n=254	NS	0	1(1.0)	3(3.1)	1(1.0)	0	0	5(5)	5(5)
	NP	0	2(1.3)	4(2.5)	1(0.6)	0	0	7(4)	7(4)
	total	0	3(1.2)	7(2.7)	2(0.8)	0	0	12(5)	12(5)

* $p < 0.05$ by chi-squared test compared with area C

3.2. Respiratory symptoms and Questionnaire analysis

Response completed 806 questionnaires were administered by parent (60% from mother). All subjects had lived in the study sites for at least 1 year. The demographic and subject characteristics were analyzed resulting in Table 6. From questionnaires analysis, children living in area A were found high prevalence of respiratory symptoms than B and C. For instance, history of allergic disease (dust and chemical) in area A, B and C were 22.8%, 11.4% and 18.5% respectively. Past illness, sinus trouble, asthmatic bronchitis and other respiratory in area A, B and C were 27.6%, 23.5% and 15.7% respectively.

Parental smoking habit had no association with respiratory symptoms of the children but smoking parent shared room with children were significantly associated with respiratory symptoms, OR= 2.01 (95% CI = 1.2-3.5). Parents in area C were found higher educated level than parents in area A and B. However, in this study, the level of education has no correlation with their children's respiratory symptoms.

The prevalence of respiratory symptoms of schoolchildren in area A, B and C were compared in Table 5. The respiratory symptoms, chronic bronchitis (CB), bronchial asthma (BA), dyspnea and wheezing (D and W), persistent cough, and persistent phlegm were higher in area A than in B and C. However, the persistent cough and persistent phlegm were significantly higher in children in area A compared with those in area C ($p < 0.05$) Table 5. The results of the logistic regression analyses are shown in Table 7. The children in area A have significantly increased prevalence of respiratory symptoms compared with those living in the area C (OR=3.41, 95% CI=1.70-6.85).

The environmental factors outside the studied school including industrial sources and traffic pollution were examined. In general, PM₁₀, benzene and 1,3 butadiene were considered as the traffic related pollutions. However, our school sites, MT (area A), KH (area B) and NS (area C) are located along the main road and close to the intersections which were commonly suffering from constant heavy traffic jams. Nevertheless, students living in area C had the least respiratory symptoms compared to those of MT (area A) and KH (area B). This finding suggested that the distance from the industrial zone accounted to the prevalence of respiratory symptoms of school children in this area.

Gender has the association with the prevalence of respiratory symptoms and interestingly prevalence of respiratory symptoms of the boys significantly higher than the girls (OR=2.17, 95% CI=1.33-3.53). Other factors such as age, residential years, home size, family members, parental smoking habits, use of air conditioners and domestic pets were not different among the areas. In this study, boys have a higher risk of the respiratory symptoms than girls 2.17 times. This could be associated to children's daily activity. Potentially, boy could enjoy outdoor rather than indoor functions, thus increasing more opportunities of pollutants exposure higher than girls. Our result is consistent to the findings of Moraes *et al.* (2010) who examined the association between wheezing in children and adolescents living downwind of dispersion plume emitted from petrochemical complex in Brazil. Even with low levels of pollutants, the prevalence of wheezing in male was higher in female 2.5 times. Liu *et al.* (2014) examined the association between outdoor and indoor air pollution and respiratory diseases among children aged 6 to 13 years living in a heavy industrial

Table 6. Demographic and risk factor characteristics of children included in the analysis, (n=806)

Parameter	Elementary student included in the analysis		
	Area A n=420	Area B n=132	Area C n=254
Mother responder, n (%)	268(63.8)	89(67.4)	150(59.0)
Boys, n(%)	165(39.3)	66(50.0)	118(46.4)
Mean age \pm SD (years)	11.0 \pm 0.9	10.8 \pm 0.8	10.6 \pm 0.9
Mean height \pm SD (cm.)	144.1 \pm 9.1	140.7 \pm 8.7	144.1 \pm 9.9
Mean weight \pm SD (kg.)	38.7 \pm 11.3	37.0 \pm 11.6	39.3 \pm 12.5
BMI \pm SD	18.35 \pm 3.83	18.43 \pm 4.15	18.57 \pm 4.32
Total born in Rayoung (%)	242(57.6)	67(50.7)	140(55.1)
Residential years, n(%)			
\leq 5	110(26.2)	53(40.2)	103(40.6)
6-10	211(50.2)	63(47.7)	109(42.9)
$>$ 10	91(21.7)	16(12.1)	42(16.5)
History of allergic diseases, n(%)			
Dust and Other chemical	96(22.8)	15(11.4)	47(18.5)
Past illness, n(%)	274(65.2)	68(51.5)	110(43.3)
Respiratory Symptoms*	116(27.6)	31(23.5)	40(15.7)
- Sinus trouble	25(6.0)	0	8(3.1)
- Asthmatic bronchitis	58(13.8)	20(15.2)	13(5.1)
- Other Respiratory	33(7.9)	11(8.3)	19(7.5)
Other			
- Measles (not German)	50(11.9)	11(8.3)	17(6.7)
- Pneumonia	19(4.5)	5(3.8)	6(2.4)
- Whooping cough	19(4.5)	3(2.3)	7(2.8)
- Bronchitis	57(13.6)	14(10.6)	34(13.4)
- Bronchiolitis	13(3.1)	4(3.0)	6(2.4)
Home size, room, n(%)			
1	154(36.7)	37(28.0)	61(24.0)
2-5	256(61.0)	91(68.9)	181(71.3)
$>$ 5	3(0.7)	1(0.7)	9(3.5)
Family members, n(%)			
1-5	340(81.0)	104(78.8)	198(78.0)
6-10	73(17.4)	22(16.7)	51(20.1)
$>$ 10	3(0.7)	1(0.8)	2(0.8)
people share his/her bedroom	89(21.2)	23(17.4)	37(14.6)
smoke cigarettes			
Cooking in home			
Gas	347(82.6)	90(68.2)	158(62.2)
Electricity	11(2.6)	5(3.8)	19(7.5)
Charcoal	13(3.1)	7(5.3)	1(0.4)
Other	48(11.4)	29(22.0)	75(29.5)
Use of air conditioners, n(%)	85(20.2)	26(19.7)	136(53.5)
Domestic pet, n(%)	232(55.2)	73(55.3)	117(46.1)
Parental smoking habits, n(%)	181(43.1)	53(40.2)	96(37.8)
Socioeconomic status			
Father' Education			
No formal education	2(0.5)	2(1.5)	0
Primary school	157(37.4)	45(34.1)	49(19.3)
High school	127(30.2)	39(29.6)	69(27.2)
\geq Bachelor's Degree	40(9.5)	12(9.1)	75(29.5)
Mother' Education			
No formal education	9(2.1)	5(3.8)	1(0.4)
Primary school	171(40.7)	47(35.6)	60(23.6)
High school	155(36.9)	46(34.9)	90(35.4)
\geq Bachelor's Degree	26(6.2)	11(8.3)	48(18.9)

*From Table 4

Table 7. Multiple logistic regression analyses for the association between some independent variables, any of the respiratory symptoms among school children in Map Ta Phut, Thailand

Independent variables	Children (n=806)
	Any of the respiratory symptoms OR (95% CI)
Gender (ref: girls)	2.17 (1.33-3.53)**
Age (ref: 9 year)	
10 year	2.36 (0.52-10.63)
11 year	1.99 (0.44-9.03)
12 year	1.74 (0.37-8.13)
Residential years (ref: 0-5 year)	
6-10 year	0.80 (0.46-1.40)
>10 year	0.85 (0.40-1.82)
Family members (ref: 1-5)	
6-10 member	1.08 (0.56-2.06)
Parental smoking habits	1.45 (0.89-2.36)
Use of air conditioners	1.00 (0.55-1.81)
Domestic pets	0.96 (0.58-1.57)
Areas ^a (ref: area C)	
A	3.41 (1.70-6.85)*
B	1.36 (0.54-3.45)

* $p < 0.001$, ** $p < 0.002$

province of China and they found that the prevalent rates of asthma and asthma related symptoms in male were significantly higher than that in female ($p < 0.05$).

4. Conclusions

Children living near industrial areas particularly with high level of PM₁₀ and VOCs are at risk of chronic bronchitis, bronchial asthma, dyspnea and wheezing, persistent cough and persistent phlegm. The results suggested a need for environmental policies to control industrial pollution and reduce the resident exposure. Further investigations regarding health effects of Cohort exposure to air pollutants in this area are recommended.

Acknowledgements

This research was funded by the National Research Council of Thailand 2014 GRB_APS_20572304 and Graduate School, Chulalongkorn University. The authors are deeply grateful to the parents and schoolchildren who participated in our study. The authors are greatly appreciate the valuable suggestion of field research design from Professor Eiji Yano, Department of Hygiene and Public Health, Teikyo University School of Medicine, Japan.

References

Aekplakorn W, Loomis D, Vichit-Vadakan N, Shy C, Plungchuchon S. Acute effects of SO₂ and particles from a power plant on respiratory symptom of children, Thailand. The Southeast Asian Journal of Tropical Medicine and Public Health 2003; 34(4): 906-14.

Air Quality and Noise Management Bureau (AQNIS), Pollution Control Department, Ministry of Natural Resources and Environment [homepage on the Internet]. Thailand [Cited 2015 January 22]. Available from: www.aqnis.pcd.go.th

ATS-DLD questionnaire [monograph on the Internet]. [Cited 2015 November 11]. Available from: <https://www.thoracic.org/statements/resources/archive/rrdquacer.pdf>

Environmental Protection Agency (EPA), AIRTrends1995 Summary [homepage on the Internet]. [Cited 2015 November 4]. Available from: <http://www3.epa.gov/airtrends/aqtrnd95/pm10.html>.

Environmental Protection Agency (EPA), [homepage on the Internet] Revised in January 2000 [Cited 2015 November 4]. Available from: http://www3.epa.gov/airtoxics/hlthef/di_ethan.html.

Environmental Protection Agency (EPA), [homepage on the Internet] Revised in January 2000; March 2009 [Cited 2015 November 4]. Available from: <http://www3.epa.gov/airtoxics/hlthef/butadien.html>.

Environmental Protection Agency (EPA), [homepage on the Internet] Revised in January 2000; January 2012 [Cited 2015 November 4]. Available from: <http://www3.epa.gov/airtoxics/hlthef/benzene.html>.

Ferris BG. Epidemiology standardization project (American Thoracic Society). The American Review of Respiratory Disease 1978; 118: 1-120.

Karita K, Yano E, Tamura K, Jinsart W. Effect of working and residential location areas on air pollution related respiratory symptoms in policemen and their wives in Bangkok, Thailand. European Journal of Public Health Association 2004; 14(1): 24-26.

Kongtip P, Singkaew P, Yoosook W, Chantanakul S, Sujiratat D. Health effects of people living close to a petrochemical industrial estate in Thailand. Journal of the Medical Association of Thailand 2013; 96(Suppl 5): S64-72.

- Langkulsen U, Jinsart W, Karita K, Yano E. Respiratory symptoms and lung function in Bangkok school children. *European Journal of Public Health* 2006; 16(6): 676-81.
- Liu F, Zhao Y, Liu YQ, Liu Y, Sun J, Huang MM, Liu Y, Dong GH. Asthma and asthma related symptoms in 23,326 Chinese children in relation to indoor and outdoor environmental factors: The Seven Northeastern Cities (SNEC) Study. *Science of the Total Environment* 2014; 497-498: 10-17.
- Moraes AC, Ignotti E, Netto PA, Jacobson Lda S, Castro H, Hacon Sde S. Wheezing in children and adolescents living next to a petrochemical plant in Rio Grande do Norte, Brazil. *Journal of Pediatrics* 2010; 86(4): 337-44.
- Map Ta Phut (MTP) Municipality, Rayong Province, [homepage on the Internet]. Thailand [Cited 2014 October 29]. Available from: <http://www.mtp.go.th>.
- Map Ta Phut Industrial Estate (MTPIE), Industrial Estate Authority of Thailand. [homepage on the Internet]. [Cited 2014 November 12]. Available from: <http://www.mtpie.com>.
- Pascal M, Pascal L, Bidondo ML, Cochet A, Sarter H, Stempfelet M, Wagner V. A review of the epidemiological methods used to investigate the health impacts of air pollution around major industrial areas. *Journal of Environmental and Public Health* 2013; 1-17.
- Peled R, Friger M, Bolotin A, Bibi H, Epstein L, Pilpel D, Scharf S. Fine particles and meteorological conditions are associated with lung function in children with asthma living near two power plants. *Public Health* 2005; 119(5): 418-25.
- Phatrabuddha N, Maharatchpong N, Keadtongtawee S, Saowakhontha S. Comparison of personal BTEX exposure and pregnancy outcomes among pregnant women residing in and near petrochemical industrial area. *EnvironmentAsia* 2013; 6(2): 34-41.
- Pimpisit D, Jinsart W, Hooper MA. Modeling of the BTX species based on an emission inventory of sources at the Map Ta Phut industrial estate in Thailand. *ScienceAsia* 2005; 31(2): 103-12.
- Pollution Control Department, Ministry of Natural Resources and Environment. Air Quality and Noise Standards [homepage on the Internet]. [Cited 2015 January 22]. Available from: http://www.pcd.go.th/info_serv/en_reg_std_airsnd01.html
- Rumchev KB, Spickett JT, Bulsara MK, Phillips MR, Stick SM. Domestic exposure to formaldehyde significantly increases the risk of asthma in young children. *The European Respiratory Journal* 2002; 20(2): 403-08.
- Rumchev KB, Spickett JT, Bulsara MK, Phillips MR, Stick SM. Association of domestic exposure to volatile organic compounds with asthma in young children. *Tharax* 2004; 59(9): 746-51.
- Tamura K, Jinsart W, Yano E, Karita K, Boudoung D. Particulate air pollution and chronic respiratory symptoms among traffic policemen in Bangkok. *Archives of Environmental Health* 2003; 58(4): 201-07.
- Tanyanont W, Vichit-Vadakan N. Exposure to volatile organic compounds and health risks among residents in an area affected by a petrochemical complex on Rayong, Thailand. *The Southeast Asian Journal of Tropical Medicine and Public Health* 2012; 43(1): 201-11.
- Thepanondh S, Varoonphan J, Sarutichart P, Makkasap T. Airborne volatile organic compounds and their potential health impact on the vicinity of petrochemical industrial complex. *Water, Air and Soil Pollution* 2011; 214(1): 83-92.
- Wichmann FA, Müller A, Busi LE, Cianni N, Massolo L, Schlink U, Porta A, Sly PD. Increased asthma and respiratory symptoms in children exposed to petrochemical pollution. *Journal of Allergy and Clinical Immunology* 2009; 123(3): 632-38.
- Wilson D, Takahashi K, Pan G, Chan CC, Zhang S, Feng Y, Hoshuyama T, Chuang KJ, Lin RT, Hwang JS. Respiratory symptoms among residents of a heavy-industry province in China: prevalence and risk factors. *Respiratory Medicine* 2008; 102(11): 1536-44.
- Yamane T. *Statistics: an introductory analysis*. 3rd ed. New York: Harper and Row. 1973.
- Yogev-Baggio T, Bibi H, Dubnov J, Or-Hen K, Carel R, Portnov BA. Who is affected more by air pollution-Sick or healthy? Some evidence from a health survey of schoolchildren living in the vicinity of a coal-fired power plant in Northern Israel. *Health and Place* 2010; 16(2): 399-408.

Received 12 October 2015

Accepted 26 November 2015

Correspondence to

Professor Wanida Jinsart

Department of Environmental Science,
Faculty of Science, Chulalongkorn University,
Phayathai Road, Bangkok 10330,
Thailand

Phone: +662 218 5181-3

Mobile: 66-81-8375127

Fax: +66 2218 5180

E-mail: jwanida@chula.ac.th